



Fibroscan Clinic

Physician Referral for Fibroscan

To order a Fibroscan for a patient please fill out all fields with an asterisk (*) and fax to:

FAX 888.713.2152

Patient Information

* Date _____

* First Name _____

*Last Name _____

MI _____

*DOB: _____

*Phone: _____

*Primary Insurance _____

Diagnosis / ICD-10 Codes

* Please identify the appropriate ICD-10 Codes from the list on the next page:

Special Instructions: _____

*Physician Phone _____

*Physician FAX _____

*Ordering Physician _____

*Physician Clinic _____

* Please check here if this patient is to be screened for enrollment in the HonorHealth NASH Clinical Trial: _____

Ordering Fibroscans

Insurance will reimburse for 2 Fibroscans per Year

There are no "approved" diagnosis codes for the Fibroscan. The clinical indications are:

Chronic Viral Diseases

- Chronic hepatitis C
- Chronic hepatitis B
- Antiviral treatment follow up
- Co infection

Metabolic Diseases

- NAFLD (non-alcoholic fatty liver disease)
- Diabetes

Alcoholic Liver Disease (ALD)

- Alcoholic Liver Disease

Portal Hypertension and Prognosis

- Portal Hypertension
- Prognosis Value

Miscellaneous

- Liver Transplantation
- Biliary Diseases

Any ICD-10 codes that correspond to the above clinical indications are acceptable. Below are sample ICD-10 codes that may apply.

Diagnosis	ICD- 10 Codes
Chronic Hepatitis C	B18.2
Chronic Hepatitis B	B18.1
NAFLD	K76.0
Alcoholic Liver Disease	K70.0
Portal Hypertension	K76.6
Liver Transplant Status	Z94.4
Abnormal Liver Function Studies	R94.5
Cirrhosis of Liver	K74.60
Hepatic Fibrosis	K74.0
Liver Disease, Unspecified	K76.9
Unspecified Jaundice	R17
Biliary Cirrhosis	K74.5
Biliuria	R82.2

Fibroscans are medically necessary for:

Patients with HCV to qualify for antiviral medications

Patients who have cleared the HCV virus with antiviral medications

Patients with NAFLD in order to screen for NASH

Patients with NASH to monitor progression or regression of fibrosis

Patients with Diabetes

ALD Patients who have achieved sobriety

Patients with elevated Liver Enzymes

Patients with Bile Duct Diseases

Liver Transplant Patients